

COMPONENTS: (1) Benzenesulfonamide, 4-amino-N-(2,6-dimethoxy-4-pyrimidinyl)- (sulfadimethoxine); $C_{12}H_{14}N_4O_4S$; [122-11-2] (2) 1-Pentanol; $C_5H_{12}O$; [71-41-0]	ORIGINAL MEASUREMENTS: Mauger, J. W.; Petersen, H., Jr.; Alexander, K. S.; Paruta, A. N. <i>Drug Dev. Ind. Pharm.</i> <u>1977</u> 3(2), 163-83.																			
VARIABLES: Temperature	PREPARED BY: R. Piekos																			
EXPERIMENTAL VALUES: <table border="1" data-bbox="319 731 924 1038" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="3">Solubility</th> </tr> <tr> <th>mg/ml</th> <th>$10^4 X^a$</th> <th>$10^3 \text{ mol dm}^{-3} b$</th> </tr> </thead> <tbody> <tr> <td>25</td> <td>0.98</td> <td>3.41</td> <td>3.16</td> </tr> <tr> <td>30</td> <td>1.26</td> <td>4.41</td> <td>4.06</td> </tr> <tr> <td>37</td> <td>1.60</td> <td>5.65</td> <td>5.16</td> </tr> </tbody> </table> <p data-bbox="381 1064 626 1091" style="margin-left: 40px;">^a X = mole fraction</p> <p data-bbox="381 1122 673 1148" style="margin-left: 40px;">^b Calculated by compiler</p>		t/°C	Solubility			mg/ml	$10^4 X^a$	$10^3 \text{ mol dm}^{-3} b$	25	0.98	3.41	3.16	30	1.26	4.41	4.06	37	1.60	5.65	5.16
t/°C	Solubility																			
	mg/ml	$10^4 X^a$	$10^3 \text{ mol dm}^{-3} b$																	
25	0.98	3.41	3.16																	
30	1.26	4.41	4.06																	
37	1.60	5.65	5.16																	
AUXILIARY INFORMATION																				
METHOD/APPARATUS/PROCEDURE: A const temp bath contg screw-capped bottles with sulfadimethoxine in excess and 1-pentanol was rotated for 24 h. Samples were withdrawn through a pledget of glass wool into a pipet, which was wiped clean and allowed to drain into a volumetric flask. Soly was detd from absorbance and previously ascertained Beer's law plots detd on a Cary model 16 spectrophotometer (1).	SOURCE AND PURITY OF MATERIALS: Sulfadimethoxine: lot 203027, Hoffmann-La Roche, Inc. Its mp agreed with the literature value. 1-Pentanol was from Fisher Scientific Co. Its refractive index value and density agreed with literature values.																			
ESTIMATED ERROR: Soly: av of at least 3 detns is reported (authors). Temp: $\pm 0.1^\circ\text{C}$ (authors).																				
REFERENCES: 1. Mauger, J. W.; Paruta, A. N.; Gerraughty, R. J. <i>J. Pharm. Sci.</i> <u>1972</u> , 61(1), 94.																				

COMPONENTS: (1) Benzenesulfonamide, 4-amino-N-(2,6-dimethoxy-4-pyrimidinyl)- (sulfadimethoxine); $C_{12}H_{14}N_4O_4S$; [122-11-2] (2) 1-Octanol; $C_8H_{18}O$; [111-87-5]	ORIGINAL MEASUREMENTS: Mauger, J. W.; Paruta, A. N.; Gerraughty, R. J. <i>J. Pharm. Sci.</i> <u>1972</u> , <i>61</i> (1), 94-7.								
VARIABLES: Temperature	PREPARED BY: R. Piekos								
EXPERIMENTAL VALUES: <table style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="text-align: center;">$t/^\circ C$</th> <th style="text-align: center;">Mole fraction solubility ($\times 10^4$)</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">25</td> <td style="text-align: center;">2.04</td> </tr> <tr> <td style="text-align: center;">30</td> <td style="text-align: center;">2.78</td> </tr> <tr> <td style="text-align: center;">37</td> <td style="text-align: center;">3.59</td> </tr> </tbody> </table>		$t/^\circ C$	Mole fraction solubility ($\times 10^4$)	25	2.04	30	2.78	37	3.59
$t/^\circ C$	Mole fraction solubility ($\times 10^4$)								
25	2.04								
30	2.78								
37	3.59								
AUXILIARY INFORMATION									
METHOD/APPARATUS/PROCEDURE: A const temp bath contg screw-capped bottles with sulfadimethoxine in excess and 1-octanol was rotated for 24 h. Samples were withdrawn through a pledget of glass wool into a pipet, which was wiped clean and allowed to drain into a volumetric flask. Solute concns were detd by spectrophotometric assay at predetd wavelengths using a Cary model 16 spectrophotometer (1).	SOURCE AND PURITY OF MATERIALS: Sulfadimethoxine: lot 203057; Hoffmann-La Roche, Inc. 1-Octanol was from Mallinckrodt Chem Works. ESTIMATED ERROR: Soly: av of 3 detns are given (authors). Temp: $\pm 0.1^\circ C$ (authors). REFERENCES: 1. Paruta, A. N.; Mauger, J. W. <i>J. Pharm. Sci.</i> <u>1971</u> , <i>60</i> , 432.								

COMPONENTS: (1) Benzenesulfonamide, 4-amino-N-(2,6-dimethoxy-4-pyrimidinyl)- (sulfadimethoxine); $C_{12}H_{14}N_4O_4S$; [122-11-2] (2) 1-Octanol; $C_8H_{18}O$; [111-87-5]	ORIGINAL MEASUREMENTS: Mauger, J. W.; Petersen, H., Jr.; Alexander, K. S.; Paruta, A. N. <i>Drug Dev. Ind. Pharm.</i> <u>1977</u> , <i>3</i> (2), 163-83.																			
VARIABLES: Temperature	PREPARED BY: R. Piekos																			
EXPERIMENTAL VALUES: <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="3">Solubility</th> </tr> <tr> <th>mg/ml</th> <th>$10^4 \times a$</th> <th>$10^3 \text{ mol dm}^{-3} b$</th> </tr> </thead> <tbody> <tr> <td>25</td> <td>0.40</td> <td>2.04</td> <td>1.29</td> </tr> <tr> <td>30</td> <td>0.54</td> <td>2.78</td> <td>1.74</td> </tr> <tr> <td>37</td> <td>0.70</td> <td>3.59</td> <td>2.26</td> </tr> </tbody> </table> <p style="margin-left: 40px;">^a X = mole fraction</p> <p style="margin-left: 40px;">^b Calculated by compiler</p>		t/°C	Solubility			mg/ml	$10^4 \times a$	$10^3 \text{ mol dm}^{-3} b$	25	0.40	2.04	1.29	30	0.54	2.78	1.74	37	0.70	3.59	2.26
t/°C	Solubility																			
	mg/ml	$10^4 \times a$	$10^3 \text{ mol dm}^{-3} b$																	
25	0.40	2.04	1.29																	
30	0.54	2.78	1.74																	
37	0.70	3.59	2.26																	
AUXILIARY INFORMATION																				
METHOD/APPARATUS/PROCEDURE: A const temp bath contg screw-capped bottles with sulfadimethoxine in excess and 1-octanol was rotated for 24 h. Samples were withdrawn through a pledget of glass wool into a pipet, which was wiped clean and allowed to drain into a volumetric flask. Soly was detd from absorbance and previously ascertained Beer's law plots detd on a Cary model 16 spectrophotometer (1).	SOURCE AND PURITY OF MATERIALS: Sulfadimethoxine: lot 203027; Hoffmann-La Roche, Inc. Its mp agreed with the literature value. 1-Octanol was from the Fisher Scientific Co. Its refractive index value and density agreed with literature values. ESTIMATED ERROR: Soly: av of at least 3 detns is reported (authors). Temp: $\pm 0.1^\circ\text{C}$ (authors). REFERENCES: 1. Mauger, J. W.; Paruta, A. N.; Gerraughty, R. J. <i>J. Pharm. Sci.</i> <u>1972</u> , <i>61</i> (1), 94.																			

COMPONENTS:	ORIGINAL MEASUREMENTS:																					
(1) Benzenesulfonamide, 4-amino-N-(2,6 -dimethoxy-4-pyrimidinyl)- (sulfadimethoxine); $C_{12}H_{14}N_4O_4S$; [122-11-2] (2) 1-Decanol; $C_{10}H_{22}O$; [112-30-1]	Mauger, J. W.; Petersen, H., Jr.; Alexander, K. S.; Paruta, A. N. <i>Drug Dev. Ind. Pharm.</i> <u>1977</u> , <i>3</i> (2), 163-83.																					
VARIABLES:	PREPARED BY:																					
Temperature	R. Piekos																					
EXPERIMENTAL VALUES:																						
<table border="1"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="3">Solubility</th> </tr> <tr> <th>mg/ml</th> <th>$10^4 X^a$</th> <th>$10^3 \text{ mol dm}^{-3} b$</th> </tr> </thead> <tbody> <tr> <td>25</td> <td>0.36</td> <td>2.24</td> <td>1.16</td> </tr> <tr> <td>30</td> <td>0.44</td> <td>2.69</td> <td>1.42</td> </tr> <tr> <td>37</td> <td>0.54</td> <td>3.37</td> <td>1.74</td> </tr> </tbody> </table>				t/°C	Solubility			mg/ml	$10^4 X^a$	$10^3 \text{ mol dm}^{-3} b$	25	0.36	2.24	1.16	30	0.44	2.69	1.42	37	0.54	3.37	1.74
t/°C	Solubility																					
	mg/ml	$10^4 X^a$	$10^3 \text{ mol dm}^{-3} b$																			
25	0.36	2.24	1.16																			
30	0.44	2.69	1.42																			
37	0.54	3.37	1.74																			
$a X = \text{mole fraction}$																						
$b \text{ Calculated by compiler}$																						
AUXILIARY INFORMATION																						
METHOD/APPARATUS/PROCEDURE:	SOURCE AND PURITY OF MATERIALS:																					
The soly was detd by the method of Paruta et al. (1): Screw-capped bottles with sulfadimethoxine in excess and 1-decanol were rotated in a const temp bath for 24 h. Samples were withdrawn through a pledget of glass wool into a pipet, which was wiped clean and allowed to drain into a volumetric flask. Soly was detd from absorbance and previously ascertained Beer's law plots detd on a Cary model 16 spectrophotometer.	Sulfadimethoxine: lot 203027, Hoffman-La Roche, Inc. M.p agreed with that of literature. 1-Decanol was purchased from Matheson, Coleman and Bell. Refractive index value and density agreed with those reported in the literature.																					
ESTIMATED ERROR:																						
Temp: $\pm 0.1^\circ\text{C}$ (authors). Soly: not specified.																						
REFERENCES:																						
1. Paruta, A. N.; Mauger, J. W.; Gerraughty, R. J. <i>J. Pharm. Sci.</i> <u>1972</u> , <i>61</i> , 94.																						