

INTRODUCTION TO THE SOLUBILITY OF LIQUIDS IN LIQUIDS

The Solubility Data Series is made up of volumes of comprehensive and critically evaluated solubility data on chemical systems in clearly defined areas. Data of suitable precision are presented on data sheets in a uniform format, preceded for each system by a critical evaluation if more than one set of data is available. In those systems where data from different sources agree sufficiently, recommended values are proposed. In other cases, values may be described as "tentative", "doubtful" or "rejected".

This volume is primarily concerned with liquid-liquid systems, but related gas-liquid and solid-liquid systems are included when it is logical and convenient to do so. Solubilities at elevated and low temperatures and at elevated pressures may be included, as it is considered inappropriate to establish artificial limits on the data presented.

For some systems the two components are miscible in all proportions at certain temperatures or pressures, and data on miscibility gap regions and upper and lower critical solution temperatures are included where appropriate and if available.

TERMINOLOGY

In this volume a mixture (1,2) or a solution (1,2) refers to a single liquid phase containing components 1 and 2, with no distinction being made between solvent and solute.

The *solubility* of a substance 1 is the relative proportion of 1 in a mixture which is saturated with respect to component 1 at a specified temperature and pressure. (The term "saturated" implies the existence of equilibrium with respect to the processes of mass transfer between phases).

QUANTITIES USED AS MEASURES OF SOLUBILITY

Mole fraction of component 1, x_1 or $x(1)$:

$$x_1 = n_1 / \sum_i n_i = \frac{m_1/M_1}{\sum_i (m_i/M_i)}$$

where n_i is the amount of substance (number of moles) of component i , m_i is the mass of substance i , and M_i is its molar mass.

Mole per cent of component 1 is $100x_1$.

Mass fraction of component 1, w_1

$$w_1 = m_1 / \sum_i m_i$$

where m_i is the mass of component i .

Mass percent of component 1 is $100w_1$, and may be described as g(1)/100 g sln which makes it clear that it is mass percent of solute relative to solution and not solvent. The equivalent terms "weight fraction" and "weight percent" are not used. The mole fraction solubility is related to the mass fraction solubility in a binary system by

$$x_1 = \frac{w_1/M_1}{w_1/M_1 + (1 - w_1)/M_2}$$

Amount-of-substance concentration of component 1 in a solution of volume V ,

$$c_1 = n_1/V$$

is expressed in units of mol dm^{-3} . The terms "molarity" and "molar" and the unit symbol M are not used.

Mass ratio is occasionally used in a two-component solution in the form $g(1)/g(2)$, $\text{mg}(1)/g(2)$, etc. The term "part per million" (ppm) is not used, but may be expressed as $\text{mg}(1)/\text{kg sln}$.

Molality of component 1 in component 2 is often used in solid-liquid systems, defined $m_1 = n_1/n_2 M_2$, with units mol kg^{-1} , but is not used in liquid-liquid systems where the distinction between "solute" 1 and "solvent" 2 is inappropriate. The term molality alone is inadequate, and the unit (mol kg^{-1} , mmol kg^{-1}) must be stated.

Mole fractions and mass fractions are appropriate to either the "mixture" or the "solution" point of view; the other quantities are appropriate to the solution point of view only.

ORDERING OF SYSTEMS

It is necessary to establish a method of ordering chemical compounds, to be used for the lists of saturating components which define each chemical system. This order is also used for ordering systems within volumes.

The systems are ordered first on the basis of empirical formula according to the Hill system (ref 2). The organic compounds within each Hill formula are ordered as follows:

- (i) by degree of unsaturation (e.g. cycloalkene, diene, alkyne, cycloalkane, alkane), then
- (ii) by order of increasing chain length in the parent hydrocarbon, then
- (iii) by order of increasing chain length of hydrocarbon branches, then
- (iv) numerically by position of unsaturation, then
- (v) numerically by position of substitution, then
- (vi) alphabetically by IUPAC name.

For example,

C_5H_8	cyclopentene 2-methyl-1,3-butadiene 1,4-pentadiene 1-pentyne
C_5H_{10}	cyclopentane 3-methyl-1-butene 2-methyl-2-butene 1-pentene 2-pentene
C_5H_{12}	2,2-dimethylpropane 2-methylbutane pentane
$C_5H_{12}O$	2,2-dimethyl-1-propanol 2-methyl-1-butanol 2-methyl-2-butanol 3-methyl-1-butanol 3-methyl-2-butanol 1-pentanol 2-pentanol 3-pentanol

$C_6H_{12}O$	cyclohexanol 4-methyl-1-penten-3-ol 1-hexen-3-ol 4-hexen-3-ol
$C_{18}H_{12}$	benzanthracene chrysene naphthacene triphenylene

Deuterated compounds immediately follow the corresponding 1H compounds.

GUIDE TO THE COMPILATIONS AND EVALUATIONS

The format used for the compilations and evaluations has been discussed in the Foreword. Additional information on the individual sections of each sheet is now provided in the following.

"Components". Each component is listed by IUPAC name (ref 1), chemical formula according to the Hill system, and Chemical Abstracts Registry Number. Also included are trivial name or names if appropriate.

"Original Measurements". References are expressed in "Chemical Abstracts" style, journal names being abbreviated, and if necessary transliterated, in the forms given by the "Chemical Abstracts Service Source Index" (CASSI).

"Variables". Ranges of variations of temperature, pressure, etc. are indicated here.

"Prepared by". The compiler is named here.

"Experimental Values". Components are described as (1) and (2), as defined in "Components". The experimental data are presented in the units used in the original paper. Thus the temperature is expressed $t/^{\circ}C$ or $T/^{\circ}F$ as in the original, and conversion to T/K is made only in the critical evaluation. However, the authors's units are expressed according to IUPAC recommendations (ref 3,4) as far as possible.

In addition, compiler-calculated values of mole fractions and/or mass percent are included if the original data do not use these units. 1975 or 1977 atomic weights (ref 5) are used in such calculations. When appropriate, conversions from concentrations to mole fractions are included in the compilation sheets, with the values and sources of the densities being quoted and referenced.

Details of smoothing equations (with limits) are included if they are present in the original publication and if the temperature or pressure ranges are wide enough to justify this procedure.

Errors in calculations, fitting equations, etc. are noted, and where possible corrected. Material inserted by the compiler is identified by the word "compiler" in parentheses.

The precision of the original data is preserved when derived quantities are calculated, if necessary by the inclusion of one additional significant figure.

"Method". An outline of the method is presented, reference being made to sources of further detail if these are cited in the original paper. "Chemical Abstracts" abbreviations are often used in this text.

"Source and Purity of Materials". For each component, referred to as (1) or (2), the following information (in this order and in abbreviated form) is provided if it is available in the original paper:

- source and specification
- method of preparation
- properties
- degree of purity.

"*Estimated Error*". If this information was omitted by the authors, and if the necessary data are available in the paper, the compilers have attempted to estimate errors (identified by "compiler" in parentheses) from the internal consistency, the type of apparatus, and other relevant information. Methods used by the compilers for reporting estimating errors are based on the papers by Ku and Eisenhart (ref 6).

"*References*". These are the references (usually cited in the original paper) which the compiler considers particularly useful in discussing the method and material.

"*Evaluator*". The information provided here is the name of the evaluator, the evaluator's affiliation, and the date of the evaluation.

"*Critical Evaluation*". The evaluator aims, to the best of his or her ability, to check that the compiled data are correct, to assess their reliability and quality, to estimate errors where necessary, and to recommend numerical values. The summary and critical review of all the data supplied by the compiler include the following information:

- (a) *Critical text*. The evaluator produces a text evaluating *all* the published data for the particular system being discussed, reviewing their merits or shortcomings. Only published data are considered, and even some of the published data may only be referred to in this text if it is considered that inclusion of a data compilation sheet is unjustified.
- (b) *Fitting equations*. If the use of a smoothing equation is justifiable, the evaluator may provide an equation representing the solubility as a function of the variables reported in the compilation sheets, stating the limits within which it should be used.
- (c) *Recommended values*. Data are *recommended* if the results of at least two independent experimental groups are available and are in good agreement, and if the evaluator has no doubt as to the adequacy and reliability of the experimental and computational procedures used.

Data are reported as *tentative* if only one set of measurements is available, or if the evaluator is uncertain of the reliability of some aspect of the experimental or computational method but judges that it should cause only minor error, or if the evaluator considers some aspect of the computational or experimental method undesirable but believes the data to have some value in those instances when an approximate value of the solubility is needed.

Data determined by an inadequate method or under ill-defined conditions is *rejected*, the reference being included in the evaluation together with a reason for its rejection by the evaluator.

(d) *References*. All pertinent references are listed here, including all those publications appearing in the accompanying compilation sheets and also those which have been rejected and not compiled.

(e) *Units*. The final recommended values are reported in SI units (ref 3).

Continuation Sheets. These are used for both compilations and evaluations, and include sections listing the "Components" and also the "Original Measurement" or "Evaluator". Compilation continuation sheets may include a section headed "Comments and/or Additional Data".

REFERENCES

1. Rigaudy, J.; Klesney, S.P. *Nomenclature of Organic Chemistry* (IUPAC), ("The Blue Book"), Pergamon, Oxford, 1979.
2. Hill, E.A. *J. Am. Chem. Soc.* 1900, *22*, 478.
3. Whiffen, D.H., ed. *Manual of Symbols and Terminology for Physicochemical Quantities and Units* (IUPAC), ("The Green Book"), Pergamon, Oxford, 1979; *Pure Appl. Chem.* 1979, *51*, 1.
4. McGlashan, M.L. *Physicochemical Quantities and Units*, 2nd ed. Royal Institute of Chemistry, London, 1971.
5. IUPAC Commission on Atomic Weight, *Pure Appl. Chem.* 1976, *47*, 75; 1979, *51*, 405.
6. Ku, H.H., and Eisenhart, C., in Ku, H.H., ed. *Precision Measurement and Calibration*, NBS Special Publication 300, Vol. 1, Washington D.C., 1969.