

COMPONENTS: (1) 3-Methyloctane; C_9H_{20} ; [2216-33-3] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Englin, B.A.; Plate, A.F.; Tugolukov, V.M.; Pryanishnikova, M.A. <i>Khim. Tekhnol. Topl. Masel</i> <u>1965</u> , 10, 42-6.												
VARIABLES: Temperature: 10-30°C	PREPARED BY: A. Maczynski and M.C. Haulait-Pirson												
EXPERIMENTAL VALUES: <p style="text-align: center;">Solubility of Water in 3-methyloctane</p> <table style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="text-align: center;">$t/^\circ C$</th> <th style="text-align: center;">$g(2)/100\ g\ sln$</th> <th style="text-align: center;">$10^4 x_2$ (compiler)</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">10</td> <td style="text-align: center;">0.0050</td> <td style="text-align: center;">3.56</td> </tr> <tr> <td style="text-align: center;">20</td> <td style="text-align: center;">0.0087</td> <td style="text-align: center;">6.20</td> </tr> <tr> <td style="text-align: center;">30</td> <td style="text-align: center;">0.0155</td> <td style="text-align: center;">11.03</td> </tr> </tbody> </table>		$t/^\circ C$	$g(2)/100\ g\ sln$	$10^4 x_2$ (compiler)	10	0.0050	3.56	20	0.0087	6.20	30	0.0155	11.03
$t/^\circ C$	$g(2)/100\ g\ sln$	$10^4 x_2$ (compiler)											
10	0.0050	3.56											
20	0.0087	6.20											
30	0.0155	11.03											
AUXILIARY INFORMATION													
METHOD/APPARATUS/PROCEDURE: Component (1) was introduced into a thermostatted flask and saturated for 5 hours with (2). Next, calcium hydride was added and the evolving hydrogen volume measured and hence the concentration of (2) in (1) was evaluated.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES:												