

| | | | |
|---|-------------------------------|--|--------------------------------|
| COMPONENTS: | | EVALUATOR: | |
| (1) 2-Methylnaphthalene; C ₁₁ H ₁₀ ; [91-57-6] | | G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia. | |
| (2) Water; H ₂ O; [7732-18-5] | | A. Maczynski, Institute of Physical Chemistry, Polish Academy of Sciences, Warszawa, Poland. | |
| | | December 1985. | |
| CRITICAL EVALUATION: | | | |
| Quantitative solubility data for 2-methylnaphthalene (1) in water (2) have been reported in the publications listed in Table 1. No data appear to have been reported on the solubility of water in 2-methylnaphthalene. | | | |
| <u>TABLE 1: Quantitative Solubility Studies of 2-Methylnaphthalene (1) in Water (2)</u> | | | |
| Reference | T/K | Method | |
| Eganhouse and Calder (ref 1) | 298 | GLC | |
| Mackay and Shiu (ref 2) | 298 | spectrofluorometric | |
| The original data in both of these publications are compiled in the Data Sheets immediately following this Critical Evaluation. | | | |
| The two values available (Table 1), obtained by different methods, are in excellent agreement and thus the mean value can be Recommended. | | | |
| <u>TABLE 2: Recommended (R) Value of the Solubility of 2-Methylnaphthalene (1) in Water (2)</u> | | | |
| T/K | Solubility values | | |
| | Reported values | "Best" value ($\pm \sigma_n$) ^a | |
| | 10 ³ g(1)/100g sln | 10 ³ g(1)/100g sln | 10 ⁶ x ₁ |
| 298 | 2.46 (ref 1), 2.54 (ref 2) | 2.50 \pm 0.04 (R) | 3.2 (R) |
| a Calculated by averaging; σ_n has no statistical significance. | | | |
| REFERENCES | | | |
| 1. Eganhouse, R.P.; Calder, J.A. <i>Geochim. Cosmochim. Acta</i> <u>1976</u> , <i>40</i> , 555-61. | | | |
| 2. Mackay, D.; Shiu, W.Y. <i>J. Chem. Eng. Data</i> <u>1977</u> , <i>22</i> , 399-402. | | | |

| | |
|---|---|
| COMPONENTS: (1) 2-Methylnaphthalene; C ₁₁ H ₁₀ ; [91-57-6] (2) Water; H ₂ O; [7732-18-5] | ORIGINAL MEASUREMENTS: Eganhouse, R.P.; Calder, J.A. <i>Geochim. Cosmochim. Acta</i> <u>1976</u> , 40, 555-61. |
| VARIABLES: One temperature: 25°C | PREPARED BY: A. Maczynski |
| EXPERIMENTAL VALUES: The solubility of 2-methylnaphthalene in water at 25°C was reported to be 24.6 mg(1)/kg(2) and 1.72×10^{-4} mol(1) dm ⁻³ (2). The corresponding mass percent and mole fraction, x_1 , calculated by the compiler are 2.46×10^{-3} g(1)/100 g sln and 3.12×10^{-6} . | |
| AUXILIARY INFORMATION | |
| METHOD/APPARATUS/PROCEDURE: A mixture of 500 mL (2) and 0.001 mol (1) was equilibrated in an Erlenmeyer flask for 12 h (agitation) + 24 h (stationary). The saturated solution, 100 mL, was extracted with hexane, concentrated by evaporation under nitrogen and analyzed by glc. A 5700 A Hewlett-Packard instrument equipped with dual compensating columns and flame ionization detectors was employed. | SOURCE AND PURITY OF MATERIALS: (1) source not specified; analytical grade; used as received; no impurities by glc. (2) doubly distilled; free of trace organics. ESTIMATED ERROR: temp. ± 0.5°C soly. ± 0.5 mg (1)/kg(2) (from eight determinations) REFERENCES: |

| | |
|---|--|
| COMPONENTS: (1) 2-Methylnaphthalene; $C_{11}H_{10}$; [91-57-6] (2) Water; H_2O ; [7732-18-5] | ORIGINAL MEASUREMENTS: Mackay, D.; Shiu, W.Y. <i>J. Chem. Eng. Data</i> <u>1977</u> , 22, 399-402. |
| VARIABLES: One temperature: 25°C | PREPARED BY: M.C. Haulait-Pirson |
| EXPERIMENTAL VALUES: The solubility of 2-methylnaphthalene in water at 25°C was reported to be 25.4 mg(1) dm^{-3} sln and $x_1 = 3.22 \times 10^{-6}$. The corresponding mass percent calculated by the compiler is 0.00254 g(1)/100 g sln. | |
| AUXILIARY INFORMATION | |
| METHOD/APPARATUS/PROCEDURE: A saturated solution of (1) in (2) was vigorously stirred in a 250 mL flask for 24 hrs. and subsequently settled at 25°C for at least 48 hrs. Then the saturated solution was decanted and filtered and 50-100 mL extracted with approximately 5 mL of cyclohexane in a separatory funnel. After shaking for 2 hrs. the cyclohexane extract was removed for analysis. An Aminco-Browman spectrophotofluorometer (American Instruments Ltd.) was used for analysis. Many details are given in the paper. | SOURCE AND PURITY OF MATERIALS: (1) Aldrich Chemicals, Eastman Kodak, or K and K Laboratories, commercial highest grade; used as received. (2) doubly distilled. ESTIMATED ERROR: soly. ± 0.2 mg(1) dm^{-3} sln (maximum deviation from several determinations). REFERENCES: |