

COMPONENTS: (1) 2,4,6-Trimethyl-2-phenylheptene; $C_{16}H_{26}$; [4810-06-4] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Englin, B.A.; Plate, A.F.; Tugolukov, V.M.; Pryanishnikova, M.A. <i>Khim. Tekhnol. Topl. Masel</i> <u>1965</u> , 10, 42-6.												
VARIABLES: Temperature: 10-30°C	PREPARED BY: A. Maczynski and Z. Maczynska												
EXPERIMENTAL VALUES: Solubility of Water in 2,4,6-Trimethyl-2-phenylheptene <table border="1" data-bbox="262 586 1044 741"> <thead> <tr> <th>$t/^\circ C$</th> <th>$g(2)/100\ g\ sln$</th> <th>$10^3\ x_2$ (compiler)</th> </tr> </thead> <tbody> <tr> <td>10</td> <td>0.0096</td> <td>1.16</td> </tr> <tr> <td>20</td> <td>0.0154</td> <td>1.87</td> </tr> <tr> <td>30</td> <td>0.0252</td> <td>3.05</td> </tr> </tbody> </table>		$t/^\circ C$	$g(2)/100\ g\ sln$	$10^3\ x_2$ (compiler)	10	0.0096	1.16	20	0.0154	1.87	30	0.0252	3.05
$t/^\circ C$	$g(2)/100\ g\ sln$	$10^3\ x_2$ (compiler)											
10	0.0096	1.16											
20	0.0154	1.87											
30	0.0252	3.05											
AUXILIARY INFORMATION													
METHOD/APPARATUS/PROCEDURE: Component (1) was introduced into a thermostatted flask and saturated for 5 hr. with (2). Next, calcium hydride was added and the evolving hydrogen volume measured and hence the concentration of (2) in (1) was evaluated.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES:												