

COMPONENTS: (1) 10-Butylbenz[a]anthracene; $C_{22}H_{20}$; [31632-63-0] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Davis, W.W.; Krahl, M.E.; Cloves, G.H.A. <i>J. Am. Chem. Soc.</i> <u>1942</u> , <i>64</i> , 108-10.												
VARIABLES: One temperature: 27°C	PREPARED BY: M.C. Haulait-Pirson												
EXPERIMENTAL VALUES: <p style="text-align: center;">Solubility of 10-butylbenz[a]anthracene in water</p> <table style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th style="text-align: center;">$t/^\circ C$</th> <th style="text-align: center;">$10^6 \text{ g(1) L}^{-1} \text{ (2)}$</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">27</td> <td style="text-align: center;">7 ± 0.7</td> </tr> <tr> <td></td> <td style="text-align: center;">7 ± 0.7</td> </tr> <tr> <td></td> <td style="text-align: center;">7 ± 0.7</td> </tr> <tr> <td></td> <td style="text-align: center;">8 ± 1</td> </tr> <tr> <td></td> <td style="text-align: center;">8 ± 1</td> </tr> </tbody> </table> <hr style="width: 50%; margin: 10px auto;"/> <p>The best value recommended by the authors is $8.0 \times 10^{-6} \text{ g(1) L}^{-1} \text{ (2)}$. With the assumption that 1.00 L sln = 1.00 kg sln, the corresponding mass percent and mole fraction, x_1, calculated by the compiler are $8.0 \times 10^{-7} \text{ g(1)/100 g sln}$ and 5.1×10^{-10}.</p>		$t/^\circ C$	$10^6 \text{ g(1) L}^{-1} \text{ (2)}$	27	7 ± 0.7		7 ± 0.7		7 ± 0.7		8 ± 1		8 ± 1
$t/^\circ C$	$10^6 \text{ g(1) L}^{-1} \text{ (2)}$												
27	7 ± 0.7												
	7 ± 0.7												
	7 ± 0.7												
	8 ± 1												
	8 ± 1												
AUXILIARY INFORMATION													
METHOD/APPARATUS/PROCEDURE: The method consisted of preparing serial dilutions of a suspension of (1) in (2) and determining nephelo- metrically the amount of (1) per unit volume beyond which further dilution caused no reduction in light scattering, which remained equal to that of pure (2). A Bausch and Lomb Dubosque colori- meter model 100-mm was employed. Many details are reported in ref 1.	SOURCE AND PURITY OF MATERIALS: (1) prepared at Harvard University; m.p. range 96.4-96.7°C. (2) dust-free. ESTIMATED ERROR: temp. ± 3°C soly. see above REFERENCES: 1. Davis, W.W.; Parker, Jr., T.V. <i>J. Am. Chem. Soc.</i> <u>1942</u> , <i>64</i> , 101. 2. Davis, W.W.; Krahl, M.E.; Cloves, G.H.A. <i>J. Am. Chem. Soc.</i> <u>1940</u> , <i>62</i> , 3086.												