

COMPONENTS: (1) Coronene; C ₂₄ H ₁₂ ; [191-07-1] (2) Water; H ₂ O; [7732-18-5]	ORIGINAL MEASUREMENTS: Mackay, D.; Shiu, W.Y. <i>J. Chem. Eng. Data</i> <u>1977</u> , 22, 399-402.
VARIABLES: One temperature: 25°C	PREPARED BY: M.C. Haulait-Pirson
EXPERIMENTAL VALUES: The solubility of coronene in water at 25°C was reported to be 0.00014 mg(1) dm ⁻³ sln and $x_1 = 8.56 \times 10^{-12}$. The corresponding mass percent calculated by the compiler is 1.4×10^{-8} g(1)/100 g sln.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: A saturated solution of (1) in (2) was vigorously stirred in a 250 mL flask for 24 hrs. and subsequently settled at 25°C for at least 48 hrs. Then the saturated solution was decanted and filtered and 50-100 mL extracted with approximately 5 mL of cyclohexane in a separatory funnel. After shaking for 2 hrs. the cyclohexane extract was removed for analysis. An Aminco-Browman spectrophotofluorometer (American Instruments Ltd.) was used for analysis. Many details are given in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Aldrich Chemicals, Eastman Kodak, or K and K Laboratories, commercial highest grade; used as received. (2) doubly distilled. ESTIMATED ERROR: soly. $\pm 2 \times 10^{-5}$ mg(1) dm ⁻³ sln (maximum deviation from several determinations). REFERENCES: