

ACKNOWLEDGEMENTS

The following figures are reprinted with the permission of the copyright owner, the American Chemical Society.

Bond, P. A.; Stephens, W. R.
J. Am. Chem. Soc. 1929, *51*, 2910-22.
 Figure 1. Miscibility of titanium tetrachloride
 with liquid sulfur dioxide. 39

Bond, P. A.; Crone, E. B.
J. Am. Chem. Soc. 1934, *56*, 2038-31.
 Figure 1. Miscibility of titanium tetrabromide
 with liquid sulfur dioxide. 57

Ferris, L. N.
J. Chem. Eng. Data 1966, *11*, 343-6.
 Figure 1. Solubility of niobium in hydrofluoric
 acid solutions at 25 °C. 234
 Figure 2. Portion of the niobium oxide-hydrofluoric
 acid-water system at 25 °C. 234

The following figure is reprinted with the permission of the copyright owner, Pergamon Press.

Larsen, E. M.; Trevorrow, I. R.
J. Inorg. Nucl. Chem. 1956, *2*, 254-9.
 Figure 2. The zirconium tetrachloride-acetonitrile-
 isoamyl ether system. 109, 177

The following figures are reprinted with the permission of the copyright owners, VAAP, The Copyright Agency of the USSR.

Tananaev, I. V.; Nikolaev, N. S.; Buslaev, Yu. A.
Zh. Neorg. Khim. 1956, *1*, 274-81.
 Figure 1. Solubility in the ZrF_4 -HF-H₂O system. 64

Nikolaev, N. S.; Buslaev, Yu. A.
Zh. Neorg. Khim. 1959, *4*, 205-12.
 Figure 3. Solubility in the HF-NbF₅-H₂O system. 227

Buslaev, Yu. A.; Nikolaev, N. S.
Zh. Neorg. Khim. 1959, *4*, 465-71.
 Figure 3. Solubility in the TaF₅-H₂O system at 20 °C. 267

Goroschenko, Ya. G.; Spasibenko, T. P.
Zh. Neorg. Khim. 1962, *7*, 1159-68.
 Figure 1. Solubility isotherms in the ZrO₂-HCl-H₂O system 118

Nikolaev, N. S.; Buslaev, Yu. A.; Gustyakova, M. P.
Zh. Neorg. Khim. 1962, *7*, 1685-92.
 Figure 1. Solubility in the HF-ZrF₄-H₂O system at 25 °C. 66

Kamaeva, I. G.; Serebrennikov, V. V.
Zh. Neorg. Khim. 1963, *8*, 2151-4.
 Figure, page 2154. Acid free projection of the ZrOCl₂ CaCl₂-
 HCl-H₂O isotherm at 25 °C. 130

Barskaya, I. B.; Morozov, I. S.
Zh. Neorg. Khim. 1968, *13*, 1408-11.
 Figure 1a-f. Solubilities at 25 °C in various systems. 1a 198
 1b 202
 1c 199
 1d 190
 1e 201
 1f 200

Figure 2b-d. Solubilities at 25 °C in various systems. 2b 191
 2c 196
 2d 197

Continued on the next page . . .

The following figures are reprinted with the permission of the copyright owners, VAAP, the Copyright Agency of the USSR.

- Baraskaya, I. B.; Toptygin, G. M.
Zh. Neorg. Khim. 1970, 15, 2572-3.
 Figures a and b. Solubilities at 25 °C in various systems. a 121
 b 203
- Blinova, Z. N.; Komissarova, L. N.
Zh. Neorg. Khim. 1972, 17, 2520-5.
 Figures 2a and b. Solubility isotherms of $\text{HfOCl}_2\text{-HCl}$
 solutions at 0 (a) and 25 °C (b). 189
- Tsurik, L. A.; Kamaeva, I. G.; Serebrennikov, V. V.
Zh. Neorg. Khim. 1977, 22, 1366-71.
 Figure 1. Acid free projection of the isotherm ($\text{HfOCl}_2\text{-CaCl}_2\text{-HCl-H}_2\text{O}$) at 25 °C. 195
- Meerson, G. A.; Zverev, G. L.; Zubkova, F. M.
Tsvetnye Metal (Non-ferrous metals) 1939, No. 8, 97-101.
 Figure 2. Phase diagram of the $\text{K}_2\text{NbOF}_5\text{-K}_2\text{TaF}_7\text{-H}_2\text{O}$ system. 275
 Figure 3. Salting out of K_2TaF_7 by K_2NbOF_5 . 283
- Delarova, N. I.; Zavaritskaya, T. A.; Zevakin, I. A.
Izv. Akad. Nauk SSSR, Otd. Tekh. Nauk, Metall. Topl.
 1960, No. 4, 33-8.
 Figure 2. Solubility of zirconium oxychloride. 40
- Buslaev, Yu. A.; Nikolaev, N. S.
Dokl. Akad. Nauk SSSR 1960, 135, 1385-7.
 Figure 1. The $\text{HF-HF}_4\text{-H}_2\text{O}$ system (isotherm at 25 °C). 153
- Buslaev, Yu. A.; Gustuakova, M. P.
Izv. Akad. Nauk. SSSR, Ser. Khim. 1963, 1533-7.
 Figure 1. Solubility in the $\text{HF-VO}_2\text{-H}_2\text{O}$ system at 25 °C. 220